

## Is It Allergies, Cold, Flu, COVID-19, or RSV?

|                           | 1000                                   |  | 12000000                                      |   |  |
|---------------------------|--|--|---|---|--|
| SYMPTOMS                  | Allergies                              | Cold   | *W<br>Flu                                     | COVID-19  | RSV  |
| Stuffy or<br>Runny Nose   | 999                                    | 999  | 9.9   | 9.9   | 000  |
| Sneezing                  | 000                                    | 0.0  | 0.0   |   | <b>©</b>   |
| Wheezing                  | 0.0                                    | 0.0  | 0.0   | 9   | 000  |
| Itchy Eyes                | 000                                    | 0  | Q.  | <b>Q</b>  |  |
| Fever                     |  | 00   | 000   | 0.0   | 00   |
| Aches & Pains             |  | 9.0  | 000   | 0.0   |  |
| Headache                  | <b>Q</b>                               | 000  | 0.0   | 00  |  |
| Cough                     | 00                                     | 999  | 000   | 000   | 000  |
| Vomiting                  |  | 0  | 0.0   | 0.0   | 00   |
| Diarrhea                  |  | Q.   | 0   | 0.0   | 99   |
| Loss of Taste<br>or Smell |  |  |   | 9.9   |  |
| Shortness of<br>Breath    |  | 0  | Q.  | 00  | 999  |
| Fatigue                   |  | 0  | 000   | 0.0   |  |
| Sore Throat               |  | 999  | 000   | 9.9   |  |
| When<br>Symptoms<br>Start | Right after<br>exposure to<br>allergen | Gets worse over<br>several days                  | Sudden start                                  | Varies  | Gets worse over<br>several days                  |
| How Long it<br>Lasts      | Weeks to Months                        | 3-14 days  | 10-14 days                                    | 10-14 days, but can last<br>months (Long COVID)   | 2-3 weeks  |
| When is it<br>Contagious? | NOT Contagious                         | 1-2 days before symptoms<br>start, then 3-4 days | 1 day before symptoms<br>start, then 3-4 days | 1-2 days before symptoms<br>start, then 5-10 days | 1-2 days before symptoms<br>start, then 3-8 days |

Sources: Nationwide Children's Hospital, nationwidechildrens.org & Free Icons, freeicons.io